



Arlington Equine  
9 Church Street  
Arlington, VT 05250  
802-375-9322



## How to Give your Horse an Intramuscular Injection

Most horse owners occasionally must give their horse an injection. Fortunately, giving an injection to a horse is an easily learned skill. Determining what type of medication the horse needs and how to administer the medication is the critical part of the process and should be determined by your veterinarian.

There are four basic types of injections used with horses.

- Intravenous injections are given into a vein.
- Intradermal injections are administered into the skin.
- Subcutaneous injections are given underneath the skin.
- Intramuscular (IM) injections are given deep into a large muscle mass where the drug is absorbed slowly into the horse's system.

Intramuscular injections are the most common type used in horses and are the focus of this handout. Although a few horses are "needle shy" and object to injections, most horses quietly accept a properly given IM injection. By following the methods outlined in this circular, horse owners should be able to safely and efficiently give an IM injection to a horse.

---

## Step 1: Safety & Preparation

### Consult Your Veterinarian

Although giving an IM injection to the horse is a routine procedure, it is not without risk to the horse. Always consult your veterinarian about the type of medication, the dosage, and the route (IM, intravenous, subcutaneous, or intradermal) before giving any drug to a horse.

Improper drug handling and injection techniques can result in infections such as needle abscesses, life-threatening drug reactions such as anaphylactic (allergic) shock, and ineffective drugs or vaccines. Have your veterinarian discuss signs of drug reaction in the horse and how to handle drug reactions.

### Precautions for Handlers

Giving an injection to a horse also can pose some risk to the human handler. A horse which seriously objects to the injection can easily injure the handler. Also, some drugs used in horses can be absorbed directly through human skin or can produce severe reactions if accidentally injected into humans by needle punctures.

## **Medication Precautions**

Before giving any injection, read the drug label on the bottle. It is a good practice to check the label before you draw the medication out of the bottle and again before you inject the drug into the horse. Check the drug name to ensure it is the one recommended by your veterinarian. Remember, the generic name for a drug and the brand name may differ.

## **Dosage**

Next check the recommended dosage. This may differ from the instructions given by your veterinarian. If you have any questions about the dosage, get clarification from your veterinarian before giving the injection.

## **Route of Injection**

Check the recommended route of injecting the drug. Again, if you have questions, check with your veterinarian.

## **Drug Handling & Storage**

Follow the recommended method of drug storage, drug handling procedures, expiration date, and precautions. Do not mix individually packaged drugs in the same injection. Compounds in one drug preparation may inactivate or decrease the efficacy of the other drug. It is better to play it safe and give the horse two separate injections.

## **Sterile Equipment**

Next, make sure you use a sterile needle and syringe. Individually packaged, disposable sterile needles and syringes are the easiest way to ensure sterile equipment. Open the packages immediately before use, and dispose of them immediately after use. Never reuse a needle because a contaminated needle can easily introduce an infection into a horse.

Never use the same needle or syringe on more than one horse. Infectious diseases can be passed easily from horse to horse this way.

## **Size of Needle**

The size of the needle depends on the medication being injected. A large-diameter needle (18 gauge) works best with thick solutions such as penicillin, while a smaller-diameter needle (20 to 21 gauge) can handle a thin, watery solution. Remember, a larger gauge number equals a smaller diameter. Larger gauge needles may break more easily than smaller gauge. If your horse reacts adversely to the injection and breaks the needle, you should make sure you can find both pieces of the needle. If you suspect a piece of needle may be retained in the horse's muscle, consult your veterinarian.

Most IM shots are given to adult horses with a 1½ -inch needle so that the medication is injected deep into the muscle mass. Foals are usually given IM injections with a 1-inch needle. Collect

used needles, with their plastic covers attached, and syringes in a resealable plastic bag or bottle and take them to your veterinarian's office for disposal in an approved medical waste container.

## **Site Preparation**

Antiseptic cleansing of the injection site is not commonly practiced by most horse owners or veterinarians. They simply brush any noticeable dirt from the injection area and insert the needle into dry skin. There is no noticeable increase in infection with this method when compared with a method that uses an antiseptic cleansing agent.

Studies have shown alcohol is a weak antiseptic that has very little effect on the cleanliness of the injection site. To thoroughly clean the injection site, the horse's hair must be shaved, the area scrubbed with an antiseptic soap which must remain in contact with the skin for at least 2 minutes, and then rinsed with alcohol. This is impractical for most horse owners and most people do not want their horse shaved at the injection site. Using a sterile needle and syringe is more important in preventing injection site infections than thoroughly cleaning the site.

## **Horse Handling**

If you are unsure how your horse will react to the injection, it is best to untie the horse and hold its lead line or have someone else hold it. A tied horse which overreacts to the injection may pull back against the tie rope. The resulting pressure on the horse's head may cause it to panic, injuring the horse or the handler. If the horse does pull back during the injection, simply move with the horse and continue the injection when it calms down. If the horse tries to kick, pull its head toward you. This automatically swings its rear end away from you. Some seriously needle-shy horses may need to be distracted by a twitch (a restraint device which tightly grips the horse's upper lip and nose) or the chain end of a stallion lead run through the mouth or over the upper gum while the injection is given.


---

## **2. Injection Site**

The site of the IM injection is important for the safety of the horse and the handler. Using a large muscle mass promotes drug absorption and decreases the chance of swelling and pain at the injection site. The site should allow the needle to be placed deep in the muscle without danger of hitting bone, ligaments, nerves, or blood vessels. Also, the injection site should allow the handler to be in a relatively safe position if the horse objects to the injection. This handout describes IM injections for the neck. Please contact our office for information on giving gluteal or pectoral injections, which can be associated with slightly higher risk to horse and handler.

### **Base of the Neck**

The base of the horse's neck is the preferred injection site. It is also favored by many horse handlers because it allows the handler to remain in a relatively safe area by the horse's shoulder. This site is a triangle defined by the nuchal ligament along the crest (top) of the horse's neck, the cervical vertebrae which form a backward S-shaped curve from the horse's poll (between its ears) toward the point of the shoulder, and the scapula (shoulder blade) (see Figure 1).



**Figure 1.** The triangle indicates the area for IM injections in the neck. Keeping near the base of neck helps prevent muscular soreness.

To locate the appropriate injection area, put the heel of your hand on the base of the horse's neck where it joins the shoulder, about midway between the crest and the bottom of the neck. The area covered by your palm is the injection site.

Higher toward the crest you risk hitting the nuchal ligaments, and lower toward the bottom of the neck is where the cervical vertebrae and blood vessels are located. Make sure you stay near the base of the neck rather than injecting higher up the neck toward the ears. This again avoids ligaments, bone, and blood vessels and gives a larger muscle mass for the injection.

---


### 3. Injection Procedure

The general procedure for an IM injection is as follows:

- remove the needle from the syringe,
- set the needle into the muscle,
- attach the syringe,
- aspirate, and
- slowly administer the medication.


Novice horse owners often worry about injecting air into the horse when the needle is inserted without the syringe attached. However this amount of air injected into the horse is minor and will not harm it.

A quiet horse may allow you to simply pop the needle into its neck or buttocks. However, if the horse needs a little distraction there are several methods of easing the needle stick. One good method is to pinch up the horse's skin next to the injection site for a few seconds prior to inserting the needle (see Figure 2). While holding the pinched skin, insert the needle next to the pinched area.



**Figure 2.** Pinching the skin near the needle injection site distracts the horse from the injection.


Another method is to hold the needle between the thumb and forefinger. Tap the horse vigorously 2 to 3 times with the side of your palm in the injection site and without breaking your rhythm, rotate your hand and insert the needle (see Figure 3). However, some horses may learn to associate the taps with the following needle stick, and leave the area prior to the needle stick.



**Figure 3.** Tapping the horse at the injection site with the side of your palm several times before inserting the needle can distract the horse from the injection.

There are several methods of giving the horse an IM injection. The method you use will depend on your experience, the injection site, and the horse's attitude. There are a few basic rules to follow no matter which injection method you use:

- 1) Insert the needle perpendicular to the skin (do not slant the needle upward or downward) and sink the needle into the muscle all the way to the hub (the place where it attaches to the syringe). This ensures that the needle will not shift during the injection and that it is deep into the muscle mass (see Figure 4).



**Figure 4.** When injecting the needle, use a quick, decisive thrust that sinks the needle in the muscle to the hub.

- 2) Always aspirate (pull back on the plunger) before injecting the drug to ensure that you are not in a blood vessel. Some commonly used drugs can kill a horse if accidentally injected into the blood stream.

If blood collects at the hub of the needle when you place it into the horse or you draw blood when you aspirate the syringe, then you must pull the needle out and redirect it before giving

the injection. It is safest to pull the needle out completely and start over in a different area with a clean needle.

However if you have a needle shy horse, you can pull the needle partially out of the skin, change the injection angle, and push it back into the muscle. Many needle shy horses object to the needle going through the skin, which is the painful part of the injection, and will be relatively quiet once the needle is through the skin. Remember to aspirate again when using a clean needle or redirecting a needle to make sure you are not in a blood vessel.

- 3) Observe the horse for any signs of allergic reaction for about 30 minutes after giving the injection. Usual signs of allergic reactions such as swelling around the injection site, hives, increased respiratory rate, depression, or agitation indicate that the horse may be allergic to the medication. Discontinue its use and consult your veterinarian immediately.

NOTE: If you are giving large, repeated doses of a medication, rotate injection sites (such as left neck, right neck, left buttock, right buttock) to reduce soreness in any one area of the horse's body. Extremely large doses (more than 15 to 20 cc) of a thick or irritating substance such as penicillin should be split into more than one injection to reduce soreness.

It is also a good practice to use more than one injection site (for example, the neck and the buttocks) when giving several separate medications or vaccines at once. Then if the horse has a drug reaction it may be easier to identify the drug that caused the problem.

---

## 4. Summary

Giving your horse an IM injection is an easily learned technique. By observing proper safety rules and injection techniques, you can often avoid infections and adverse drug reactions. A summary of the IM injection procedure follows.

1. Consult your veterinarian about the type of drug needed, route of administration, dosage, drug handling precautions, and adverse drug reactions.
2. Read the drug label.
3. Use only sterile needles and syringes.
4. Untie the horse if you are not sure of its reaction.
5. Insert the needle straight into the muscle and up to the hub.
6. Attach the syringe to the needle.
7. Aspirate (pull back) on the plunger. If blood appears in the syringe, remove the needle and try again with a clean one.
8. Slowly inject the medication.
9. Observe the horse for signs of adverse drug reaction. In some cases, you may have epinephrine ready for injection in case of anaphylactic shock.
10. Properly dispose of your needle and syringe in a medical waste container.